Contents

SMEDLEY, M. J. and STANISSTREET, M. Calcium and neurulation in mammalian embryos	1-14
sмітн, L. J. Embryonic axis orientation in the mouse and its correlation with blastocyst relationships to the uterus. II. Relationships from 4½ to 9½ days	15-35
KINTNER, C. R. and BROCKES, J. P. Monoclonal antibodies to the cells of a regenerating limb	37–55
SALÓ, E. and BAGUÑÀ, J. Cell movement in intact and regenerating planarians. Quantitation using chromosomal, nuclear and cytoplasmic markers	57-70
MEULER, D. C. and MALACINSKI, G. M. An analysis of protein synthesis patterns during early embryogenesis of the urodele – Ambystoma mexicanum	71-92
OSTER, G. F., MUREAY, J. D. and MAINI, P. K. A model for chondrogenic condensations in the developing limb: the role of extracellular matrix and cell tractions	93-112
SÁNCHEZ, E. R. and ERICKSON, R. P. Expression of the Tcp-1 locus of the mouse during early embryogenesis	113-122
sánchez, E. R., Hammerberg, C. and Erickson, R. P. Quantitation of two-dimensional gel proteins reveals unequal amounts of <i>Tcp-1</i> gene products during mouse spermatogenesis but no correlation with transmission ratio distortion	123-131
SMITH, R. K. W. and JOHNSON, M. H. DNA replication and compaction in the cleaving embryo of the mouse	133–148
MILLER, J. G. O. and SCHULTZ, G. A. Amino acid transport in mouse blastocyst compartments	149-158
DUNLOP, S. A. and BEAZLEY, L. D. Cell distributions in the retinal ganglion cell layer of adult Leptodactylid frogs after premetamorphic eye rotation	159-173
FLEMING, T. P. and PICKERING, S. J. Maturation and polarization of the endocytotic system in outside blastomeres during mouse preimplantation development	175-208
LAING, N. G. and LAMB, A. H. Muscle fibre types and innervation of the chick embryo limb following cervical spinal cord removal	209-222
KIDDER, G. M. and CONLON, R. A. Utilization of cytoplasmic poly(A) ⁺ RNA for protein synthesis in preimplantation mouse embryos	223-234
KHANER, O., MITRANI, E. and EYAL-GILADI, H. Developmental potencies of area opaca and marginal zone areas of early chick blastoderms	235-241
KIMATA, K., SAKAKURA, T., INAGUMA, Y., KATO, M. and NISHIZUKA, Y. Participation of two different mesenchymes in the developing mouse mammary gland: synthesis of basement membrane components by fat pad precursor cells	243-257
NEFF, A. W., MALACINSKI, G. M. and CHUNG, HM. Microgravity simulation as a probe for understanding early Xenopus pattern specification	259-274
JEFFERY, w. R. Identification of proteins and mRNAs in isolated yellow crescents of ascidian eggs	275-287
DALE, L., SMITH, J. C. and SLACK, J. M. W. Mesoderm induction in Xenopus laevis: a quantitative study using a cell lineage label and tissue-specific antibodies	289-312
TOMLINSON, A. The cellular dynamics of pattern formation in the eye of Drosophila	313-331
WIGMORE, P. and HOLDER, N. Regeneration from isolated half limbs in the upper arm of the axolotl	333-347
INGHAM, P. w. A clonal analysis of the requirement for the trithorax gene in the diversification of segments in Drosophila	349-365
GAY, S. W. and KOSHER, R. A. Prostaglandin synthesis during the course of limb cartilage differentiation in vitro	367-382
TAYLOR, J. S. H. and GAZE, R. M. The effects of the fibre environment on the paths taken by regenerating optic nerve fibres in <i>Xenopus</i>	383-401
INDEX OF AUTHORS AND TITLES	403-404
SUBJECT INDEX	405-409